

## New Eclipsing Binary Systems: HD 189735 and HD 116274

### M.D. Koppelman

*Dept. of Astronomy, University of Minnesota, 116 Church St. S. E., Minneapolis, MN 55455*

### Patrick Wils

*Vereniging Voor Sterrenkunde, Aarschotse Baan 31, B-3191 Boortmeerbeek, Belgium*

*Presented at the 92nd Spring Meeting of the AAVSO, April 26, 2003*

**Abstract** We present two new eclipsing binary stars: HD 189735 (EB,  $P = 0.78154d$ ) and HD 116274 (EW,  $P = 0.60471d$ ). Light curves, periods, classifications, and Wilson-Devinney models are presented and discussed. Results for HD 189735 were originally presented at the Spring Meeting of the AAVSO in Tucson, Arizona in April of 2003.

### 1. Discussion

Variability of HD 189735 was identified by WS statistics (Welch and Stetson 1993) in data acquired by the Amateur Sky Survey (TASS: Droege (2002), and Henden (2001)). Variability of HD 116274 was discovered in data acquired by Stardial (McCullough and Thakkar 1997) with the procedure described in Wils (2003). Extensive followup observations were performed to characterize the light curves, in  $R_c$  for HD 189735 and in  $VR_c$  for HD 116274. TASS  $I_c$  data are not presented because they appear to have night-to-night zeropoint problems. Data were taken through standard filters, but not transformed to the standard system. Zeropoints for the photometry were obtained by using transformed Tycho-2 magnitudes (Hog 2000) for the ensemble comparison stars.

We modeled the stars using the current Wilson-Devinney code (wd; Wilson and Devinney 1971; Wilson 1979). For HD 189735 the wd code was run in mode 2 with an assumed temperature of  $T_1 = 9000K$  for the primary, based on the MK calibration of the HD spectral type of A2. Mass ratios in the range  $0.2 < q < 0.4$  all gave equivalent results. The adjusted parameters are shown in Table 1. For HD 116274 the wd code was run in mode 3 with an assumed temperature for the primary of  $T_1 = 7200K$  based on the HD spectral type of F0. The system has an apparent total eclipse, which yields a highly accurate estimate of the mass ratio. The adjusted parameters for HD 116274 are shown in Table 2. The light curves, wd fits, and residuals are shown in Figures 1 and 2.

The ephemerides of the systems are:

$$\text{HD 189735: HJD}_{\text{MinI}} = 2452458.847 + 0.78154E$$

$$\text{HD 116274: HJD}_{\text{MinI}} = 2452772.717 + 0.60471E$$

In the manner of Terrell and Henden (2002) we describe HD 189735 as a short-period Algol eclipsing binary of type EB. HD 116274 is clearly an EW-type W UMa over-contact eclipsing binary system. It is unclear whether the apparent increased scatter at the minima of HD 116274 is observational error or whether it reflects the actual behavior of the system. The  $1-\sigma$  of the comparison star magnitudes was  $< 0.02$  magnitude for all data presented.

## 2. Acknowledgements

Many thanks are due to Tom Droege, Dirk Terrell, Arne Henden, John Greaves, and everyone else involved with The Amateur Sky Survey. Professor R. Wilson (University of Florida) is acknowledged for providing the Wilson-Devinney code. This research made use of the SIMBAD database, operated by the Centre de Données Astronomiques de Strasbourg. The  $R_c$  filter used by Koppelman is courtesy of the AAVSO. Also thanks to C. W. Robertson, Aaron Price, Lou Cohen, and Mike Linnolt for additional observations of HD 189735.

## References

- Droege, T. 2003, The Amateur Sky Survey (TASS) <http://www.tass-survey.org/>.  
Henden, A. A. 2001, *J. Amer. Assoc. Var. Star Observers*, **29**, 118.  
Hog, E., Fabricius, C., Makarov, V. V., Urban, S., Corbin, T., Wycoff, G., Bastian, U., Schwekendiek, P., and Wicenec, A. 2000, *Astron. Astrophys.*, **355**, L27.  
Mc Cullough, P., and Thakkar, U. 1997, *Publ. Astron. Soc. Pacific*, **109**, 1264.  
Terrell, D., and Henden, A. A. 2002, *Inf. Bull. Var. Stars*, No. 5310.  
Welch, D. L., and Stetson, P. B. 1993, *Astron. J.*, **105**, 1813.  
Wils, P. 2003, *Inf. Bull. Var. Stars*, No. 5401.  
Wilson, R. E. and Devinney, E. J. (WD) 1971, *Astrophys. J.*, **166**, 605.  
Wilson, R. E. 1979, *Astrophys. J.*, **234**, 1054.

Table 1. Adjusted Parameters of Light Curve Solution for HD 189735.

<i>Parameter</i>	<i>Value</i>	<i>Standard Error</i>
<i>i</i>	71°5	0°2
$T_1$	9000 K	(assumed)
$T_2$	5050 K	80 K
<i>q</i>	0.25	0.02
$\Omega_1$	2.83	0.04
$L_1/(L_1+L_2)_V$	0.96	0.01
$L_1/(L_1+L_2)_R$	0.94	0.01
$HJD_0$	2452458.847	0.0011
<i>P</i>	0 <sup>d</sup> :78154	0 <sup>d</sup> :000002

Table 2. Adjusted Parameters for Light Curve Solution for HD 116274.

<i>Parameter</i>	<i>Value</i>	<i>Standard Error</i>
<i>i</i>	82°7	0°6
$T_1$	7200 K	(assumed)
$T_2$	7300 K	80 K
<i>q</i>	0.235	0.002
$\Omega_1, \Omega_2$	2.289	0.004
$L_1/(L_1+L_2)_V$	0.79	0.01
$L_1/(L_1+L_2)_R$	0.79	0.01
$HJD_0$	2452772.717	0.0011
<i>P</i>	0 <sup>d</sup> :60471	0 <sup>d</sup> :00001




Figure 1. HD 189735. TASS *V* (filled circles) and Koppelman  $R_c$  (triangles).




Figure 2. HD 116274. Koppelman *V* (triangles) and  $R_c$  (circles).